



AGOG Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-59515

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

LEAFLET

AGOPHENOL SUSPENSION

(Chloramphenicol Oral Suspension BP)

Drug Category :
Antibiotic

Generic Name :
Chloramphenicol Palmitate.

Contents :
Susp.: Each 5 ml contains Chloramphenicol palmitate BP equivalent to 125 mg Chloramphenicol base.

Indications :
Typhoid fever, H. influenzae meningitis, severe infections, when careful clinical assessment indicates no other antibiotics are effective.

Dosage :
Adults : 50 mg / kg daily in divided doses six hourly.
Children : Under 2 weeks half adult dose ; over 2 weeks, same as adult.
or as directed by the physician.

Contra-indications :
Pregnancy, breast - feeding, porphyria, History of previous hypersensitivity to and / or toxicity from Chloramphenicol.

Precautions :
Avoid repeated courses and prolonged treatment, reduce dose in hepatic / renal impairment; blood counts required before and periodically during treatment; monitor plasma - Chloramphenicol concentrations in neonates. Serious and fatal blood dyscrasias are known to occur after the administrations of Chloramphenicol



AGOG Pharma Ltd.

(WHO - GMP CERTIFIED - GOVT RECOGNISED EXPORT HOUSE)



CIN: 11-09919

Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208. INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax: 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

1.5.3 Patient information leaflet (PIL)

Blood dyscrasias have occurred after both short term and prolonged therapy. Chloramphenicol must not be used when less potentially dangerous agents are effective. It must not be used in treatment of trivial infections, infections other than those indicated or as a prophylactic agent to prevent bacterial infection.

Interactions :

Rifampicin, Warfarin, sulphonylureas, Phenobarbitol, Phenytoin, Cyclosporin, Lactulimus.

Adverse effects :

Blood disorders including reversible and irreversible aplastic anaemia (with reports of resulting leukaemia), peripheral neuritis, optic neuritis, erythema multiforme, nausea, vomiting, glossitis, stomatitis, diarrhoea, nocturnal haemoglobinuria reported, grey syndrome (abdominal distension, pallid cyanosis, circulatory collapse) may follow excessive doses in neonates with immature hepatic metabolism.

Storage Condition :

Store under normal storage conditions (15°C-30°C).

Protect from light.

Keep all medicines out of reach of children.

Shake well before use.

Presentation :

60ml / 100 ml in amber coloured pet bottles.

Manufactured in India by



AGOG PHARMA LTD.

Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd.,
Vasai (E), Dist. Thane. INDIA