

**1.4 Product Information****1.4.3 Patient information leaflet****FLAIROF (Roflumilast Tablets 250 mcg)**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Roflumilast Tablets 250 mcg is and what it is used for
2. What you need to know before you take Roflumilast Tablets 250 mcg
3. How to take Roflumilast Tablets 250 mcg
4. Possible side effects
5. How to store Roflumilast Tablets 250 mcg
6. Contents of the pack and other information

1. What Roflumilast Tablets 250 mcg is and what it is used for

Roflumilast Tablets 250 mcg contains the active substance roflumilast, which is an anti-inflammatory medicine called phosphodiesterase-4 inhibitor. Roflumilast reduces the activity of phosphodiesterase-4, a protein occurring naturally in body cells. When the activity of this protein is reduced, there is less inflammation in the lungs. This helps to stop narrowing of airways occurring in **chronic obstructive pulmonary disease (COPD)**. Thus Roflumilast Tablets 250 mcg eases breathing problems.

Roflumilast Tablets 250 mcg is used for maintenance treatment of severe COPD in adults who in the past had frequent worsening of their COPD symptoms (so-called exacerbations) and who have chronic bronchitis. COPD is a chronic disease of the lungs that results in tightening of the airways (obstruction) and swelling and irritation of the walls of the small air passages (inflammation). This leads to symptoms such as coughing, wheezing, chest tightness or difficulty in breathing. Roflumilast Tablets 250 mcg is to be used in addition to bronchodilators.

**2. What you need to know before you take Roflumilast Tablets 250 mcg****Do not take Roflumilast Tablets 250 mcg**

- if you are allergic to roflumilast or any of the other ingredients of this medicine (listed in section 6)
- if you have moderate or severe liver problems.

Warnings and precautions

Talk to your doctor or pharmacist before taking Roflumilast Tablets 250 mcg.

Sudden attack of breathlessness

Roflumilast Tablets 250 mcg is not intended for the treatment of a sudden attack of breathlessness (acute bronchospasms). In order to relieve a sudden attack of breathlessness it is very important that your doctor provides you with another medicine to be available to you at all times that can cope with such an attack. Roflumilast Tablets 250 mcg will not help you in this situation.

Body weight

You should check your body weight on a regular basis. Talk to your doctor if, while taking this medicine, you observe an unintentional loss of body weight (not related to a diet or exercise programme).

Other diseases

Roflumilast Tablets 250 mcg is not recommended if you have one or more of the following diseases:

- severe immunological diseases such as HIV infection, multiple sclerosis (MS), lupus erythematosus (LE) or progressive multifocal leukoencephalopathy (PML)
- severe acute infectious diseases such as acute hepatitis
- cancer (except basal-cell carcinoma, a slow-growing type of skin cancer)
- or severe impairment of the heart function

There is a lack of relevant experience with Roflumilast Tablets 250 mcg under these conditions. You should talk to your doctor, if you are diagnosed with any of these diseases.

Experience is also limited in patients with a previous diagnosis of tuberculosis, viral hepatitis, herpes viral infection or herpes zoster. Please talk to your doctor if you have one of these diseases.

**Symptoms you should be aware of**

You may experience diarrhoea, nausea, abdominal pain or headache during the first weeks of treatment with Roflumilast Tablets 250 mcg. Talk to your doctor if these side effects do not resolve within the first weeks of treatment.

Roflumilast Tablets 250 mcg is not recommended in patients with a history of depression associated with suicidal thinking or behaviour. You may also experience sleeplessness, anxiety, nervousness, or depressive mood. Before starting treatment with Roflumilast Tablets 250 mcg, inform your doctor if you are suffering from any symptoms of this kind and of any additional medicines you may take since some of those could increase the probability of these side effects. You or your caregiver should also immediately inform your doctor of any changes in behaviour or mood and of any suicidal thoughts you may have.

Children and adolescents

Do not give this medicine to children and adolescents under 18 years of age.

Other medicines and Roflumilast Tablets 250 mcg

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, especially the following:

- a medicine containing theophylline (a medicine to treat respiratory diseases), or
- a medicine used for treatment of immunological diseases, such as methotrexate, azathioprine, infliximab, etanercept, or oral corticosteroids to be taken long-term.
- a medicine containing fluvoxamine (a medicine to treat anxiety disorders and depression), enoxacin (a medicine to treat bacterial infections) or cimetidine (a medicine to treat stomach ulcers or heartburn).

The effect of Roflumilast Tablets 250 mcg may be reduced if taken together with rifampicin (an antibiotic medicine) or with phenobarbital, carbamazepine or phenytoin (medicines usually prescribed for the treatment of epilepsy). Ask your doctor for advice.

Roflumilast Tablets 250 mcg may be taken with other medicines used in the treatment of COPD such as inhaled or oral corticosteroids or bronchodilators. Do not stop taking these medicines or reduce their dose unless advised by your doctor.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. You should not become pregnant during treatment with this medicine and should use an effective method of contraception during therapy, because Roflumilast Tablets 250 mcg may be harmful for the



unborn baby.

Driving and using machines

Roflumilast Tablets 250 mcg has no influence on the ability to drive and use machines.

Roflumilast Tablets 250 mcg contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Roflumilast Tablets 250 mcg

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- **For the first 28 days** - the recommended starting dose is one 250 micrograms tablet once daily.
 - The starting dose is a low dose used to help your body get used to the medicine before you start taking the full dose. At this low dose you will not get the full effect from the medicine – therefore it is important that you move on to the full dose (called a ‘maintenance dose’) after 28 days.
- **After 28 days** - the recommended maintenance dose is one 500 micrograms tablet once daily.

Swallow the tablet with some water. You may take this medicine with or without food. Take the tablet at the same time every day.

You may need to take Roflumilast Tablets 250 mcg for several weeks to achieve its beneficial effect.

If you take more Roflumilast Tablets 250 mcg than you should

If you have taken more tablets than you should, you may experience the following symptoms: headache, nausea, diarrhoea, dizziness, throbbing of your heart, light-headedness, clamminess and low blood pressure. Tell your doctor or pharmacist straight away. If possible take your medicine and this leaflet with you.

If you forget to take Roflumilast Tablets 250 mcg

If you forget to take a tablet at the usual time, take the tablet as soon as you remember on the same day. If on one day you have forgotten to take a tablet of Roflumilast Tablets 250 mcg, just carry on the next day with the next tablet as usual. Continue taking your medicine at the usual times. Do not take a double dose to make up for a forgotten dose.

If you stop taking Roflumilast Tablets 250 mcg

It is important to continue taking Roflumilast Tablets 250 mcg for as long as prescribed by



your doctor, even when you have no symptoms, in order to maintain control of your lung function.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

You may experience diarrhoea, nausea, stomach ache or headache during the first weeks of treatment with Roflumilast Tablets 250 mcg. Talk to your doctor if these side effects do not resolve within the first weeks of treatment.

Some side effects could be serious. In clinical studies and post-marketing experience, rare instances of suicidal thinking and behaviour (including suicide) were reported. Please notify your doctor immediately of any suicidal thoughts you may have. You may also experience sleeplessness (common), anxiety (uncommon), nervousness (rare), panic attack (rare) or depressive mood (rare).

In uncommon cases allergic reactions may occur. Allergic reactions may affect the skin and in rare cases cause swelling of the eyelids, face, lips and tongue, possibly leading to difficulties in breathing and/or a drop in blood pressure and accelerated heartbeat. In case of an allergic reaction, stop taking Roflumilast Tablets 250 mcg and contact your doctor immediately, or go immediately to the emergency department in the nearest hospital. Take all your medicines and this leaflet with you and provide full information of your current medications.

Other side effects include the following:

Common side effects (may affect up to 1 in 10 people)

- diarrhoea, nausea, stomach ache
- weight decrease, decreased appetite
- headache

Uncommon side effects (may affect up to 1 in 100 people)

- trembling, sensation of spinning head (vertigo), dizziness
- sensation of rapid or irregular heartbeat (palpitations)
- gastritis, vomiting
- reflux of stomach acid to the gullet (acid regurgitations), indigestion
- rash
- muscle pain, muscle weakness or cramps
- back pain
- feeling of weakness or tiredness, feeling unwell.

**Module 1-Administrative and Regional Information****Rare side effects (may affect up to 1 in 1,000 people)**

- male breast enlargement
- decreased sense of taste
- respiratory tract infections (excluding pneumonia)
- bloody stools, constipation
- elevation of liver or muscle enzymes (seen in blood tests)
- wheals (urticaria).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Roflumilast Tablets 250 mcg

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information**What Roflumilast Tablets 250 mcg contains**

The active substance is roflumilast.

Each Roflumilast Tablets 250 mcg tablet contains 250 micrograms roflumilast.

The other ingredients are Mannitol, microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate.

What Roflumilast Tablets 250 mcg looks like and contents of the pack

Roflumilast Tablets 250 mcg are White to off white, round, flat bevel edged tablets debossed with 'H' on one side and 'T' on the other side.

Each pack contains 15 tablets.

FLAIROF (Roflumilast Tablets 250 mcg)



Module 1-Administrative and Regional Information

Marketing Authorisation Holder

(Company) Name : Hetero Labs Limited,

Address : 7-2-A2, Hetero Corporate, Industrial Estate, Sanath Nagar,
Hyderabad, Telangana, Pin Code - 500018

Country : INDIA

Telephone : +91-40-23704923/24/25

Tele fax : +91-40-23704926/23714250

Manufacturer

Name : Hetero Labs Limited, Unit III,

Address : 22-110, IDA, Jeedimetla, Hyderabad, Telangana,

Country : INDIA

Telephone No.: +91-40-23096171/172/173/174

Fax No. : +91-40-23095105.