

Patient Information Leaflet
Renechlor Capsules
Chloramphenicol BP 250mg

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or your pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

What is in this leaflet:

- a. What Renechlor capsules are and what they are used for
- b. What you need to know before you take Renechlor capsules
- c. How to take Renechlor capsules
- d. Possible side effects
- e. How to store Renechlor capsules
- f. Further information

a. What Renechlor capsules are and what they are used for

Renechlor capsules contain chloramphenicol which is an antibiotic for specific uses.

They are used for treatment of typhoid fever and other dangerous infections where other antibiotics are not effective.

b. What you need to know before you take Renechlor capsules

Do not take Renechlor capsules:

- if you are allergic to chloramphenicol or any of the other ingredients (listed in section f)
- if you have recently been vaccinated
- if you have porphyria (a metabolism disorder which causes abdominal pains and mental disorder)
- if you have a minor infection
- if you are taking medicines that may suppress your bone marrow
- if you are pregnant or breast-feeding.

If any of the above apply to you, speak to your doctor or pharmacist.

Warnings and precautions

Talk to your doctor or pharmacist before taking Renechlor capsules

- if you have liver or kidney problems
- if you are elderly
- if you are taking any of the medicines listed in the next section “Other medicines and Renechlor capsules”.

Children

Do not give this medicine to children because it is unlikely to be safe.

Other medicines and Renechlor capsules

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicines may be affected by chloramphenicol or they may affect how well chloramphenicol will work.

Tell your doctor or pharmacist if you are taking:

- paracetamol
- medicines that thin the blood (e.g. warfarin)
- antidiabetics (sulphonylureas such as gliclazide)
- medicines that may suppress the bone marrow (e.g. carbamazepine, penicillamine, sulphonamides, phenylbutazone)
- medicines to treat cancer
- antipsychotic drugs (e.g. clozapine, flupenthixol, fluphenazine, procainamide)
- sulphonamide antibiotics (e.g. co-trimoxazole)
- anticonvulsants for epilepsy or related conditions (e.g. phenytoin)
- barbiturates (e.g. phenobarbitone)
- immunosuppressants (e.g. ciclosporin or tacrolimus)
- penicillin or rifampicin type antibiotics
- vitamin B12 (hydroxocobalamin)
- oral contraceptives containing oestrogen (the pill)
- medicines that treat over-active thyroid disease (e.g. carbimazole, propylthiouracil)
- anti-retroviral medicines that treat HIV infection (e.g. zidovudine).

Pregnancy, breast-feeding and fertility

Do not take Renechlor capsules if you are pregnant, planning to become pregnant or breast-feeding as they may harm the baby.

Chloramphenicol may cause a condition called Grey Baby Syndrome, where the baby's skin appears grey and the baby is listless and weak. This may be fatal and requires immediate medical attention.

Ask your doctor, pharmacist or nurse for advice before taking any medicine.

Driving and using machines

Renechlor capsules have no or negligible influence on the ability to drive and use machines.

c. How to take Renechlor capsules

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Swallow the capsules whole with a drink of water.

Adults and elderly

The usual dose is 500 mg (2 capsules) every 6 hours or 50 mg per kg body weight daily in divided doses.

Blood tests may be carried out to check the level of chloramphenicol in your body and to make sure your blood is functioning properly, as chloramphenicol can damage blood cells.

Use in children

This medicine is not suitable for children.

If you take more Renechlor capsules than you should

It is likely that you will be taking these capsules in hospital and, therefore, it is unlikely that you will overdose.

However, if you do take more capsules than you should:

1. Tell your doctor, pharmacist, nurse or nearest hospital casualty department immediately.
2. Take the container and any remaining capsules with you so that people can see what you have taken.
3. Do this even if you feel well.

If you forget to take Renechlor capsules

If you forget to take a dose take it as soon as you remember, but if it is almost time for your next dose, skip the missed dose and continue as usual.

Do not take a double dose to make up for a forgotten dose.

Tell the doctor if you miss a dose.

If you stop taking Renechlor capsules

Do not stop just because you feel better. If you stop too soon, the infection may come back. Keep taking your capsules until the prescribed course is finished.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

d. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking the capsules and **seek medical help immediately** if you have any of the following **allergic reactions**:

- difficulty breathing or swallowing, swelling of the face, lips, tongue or throat
- severe itching of the skin, with a red rash or raised lumps
- blistering of the mouth, eyes, or genital region, patchy areas of rash, peeling skin.

Seek immediate medical attention if you have any of the following symptoms:

- drowsiness, feeling faint, sudden loss of consciousness, shortness of breath, nausea, vomiting
- unusual or increased bleeding or bruising
- fever, sore throat, mouth ulcers, repeated infections or infections that will not go away
- feeling tired, breathless, and looking pale
- sudden loss or blurring of vision, loss of colour vision, eye pain which worsens with eye movement
- ringing in the ears or temporary hearing loss, head-spins (vertigo), dizziness.

The side effects listed below have been reported: Not known: frequency cannot be estimated from the available data

- inflamed, sore and swollen tongue, sore mouth, difficulty chewing and eating
- fever, nausea, vomiting, diarrhoea, abdominal pain and/or bloating, blood in stool, fatigue, dehydration, swelling.

Use of chloramphenicol in **pregnancy** and in **newborn babies** can result in Grey Baby Syndrome, where the baby's skin appears grey and the baby is listless and weak. This may be fatal and requires immediate medical attention.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine.

e. How to store Renechlor capsules

Do not store above 30°C.

Protect from light.

Keep out of reach of children.

Do not take after the expiry date which is stated on the label and on the carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

f. Further information

What Renechlor capsules contain

The active substance is chloramphenicol (250 mg).

The other ingredients are Maize starch, Colloidal silicon dioxide, Purified talc, Magnesium stearate

What Renechlor capsules look like and contents of the pack

Off white coloured size '2' capsules with "Chloro" and "250" imprinted on body and cap respectively in black ink containing off white colour powder.

Aluminium/PVC blister strip of 10 capsules and 10 of such blister strips are packed in a unit box with pack insert.

MANUFACTURED BY
RENE INDUSTRIES LIMITED
Plot 680, Kamuli, Kireka
P.O Box 6034, Kampala, Uganda



A milestone to Golden Health