

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 28 of 62

1.5 Product Information

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

VERZOL® tablet

VERZOL® oral suspension 4%

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Tablet

Each tablet contains albendazole 400 mg.

Oral suspension

10 ml of suspension contain albendazole 400 mg.

Excipients:

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tablet.

Light orange coloured, elongated, biconvex scored tablet.

Oral suspension

Orange coloured suspension.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Intestinal and skin infections

- Threadworms (*Enterobius vermicularis*),
- Roundworms (*Ascaris lumbricoïdes*),
- Hookworms (*Ankylostoma duodenale*, *Necator americanus*),
- Whipworms (*Trichuris trichuria*),
- Anguillulosis (*Strongyloides stercoralis*),
- Taeniasis (*Taenia saginata*, *Taenia solium*),
- Giardiasis (*Gardia intestinalis* or *duodenalis*) in children,

Systemic infections

- Trichinosis (*Trichinella spiralis*).

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 29 of 62

4.2 Posology and method of administration

Posology

Indications	Daily dose	Treatment duration
Intestinal and skin infections (short-term treatment with lower dose)		
Oxyurosis	Children from 1 to 2 years: 5 ml suspension (200 mg) in one single dose Adults and children older than 2 years*: 400 mg, 1 single tablet or 10 ml of suspension in single dose Strict hygiene measures should be taken and family environment should also be treated.	Single dose to be repeated 7 days after.
Roundworms Hookworms Whipworms	Children from 1 to 2 years: 5 ml of suspension (200 mg) Adults and children older than 2 years*: 400 mg, 1 single tablet or 10 ml of suspension in single dose.	Single dose. **
Anguillulosis Taeniasis (associated with others parasitosis)	Adults and children older than 2 years *: 400 mg, 1 tablet or 10 ml of suspension daily	1 daily dose during 3 days. **
Giardiasis	Children older than 2 years*: 1 tablet or 10 ml of suspension daily.	1 daily dose during 5 days.
Systemic infections (long-term treatment with higher doses)		
Trichinosis	Children*: 15 mg/kg/day divided into two daily doses Adults: 1 tablet or 10 ml of suspension twice daily	2 daily doses (morning & evening) during 10 to 15 days depending on the severity of the symptoms and on the onset of treatment.

* For children under 6 years, tablet form of 400 mg is inappropriate due to wrong route risk, and only suspension form should be used.

**If the worm control performed 3 weeks after the treatment is positive, a second treatment should be administered.

Special populations

Elderly people:

Data concerning patients from 65 years old are limited. Reports suggest that no adaptation of the posology is required in elderly people. However, albendazole should be used with care in patients with a liver dysfunction.

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 30 of 62

Liver failure:

Albendazole is rapidly metabolized by the liver, the main metabolite, albendazole sulfoxide, is pharmacologically active. Hence, liver failure might result in significant effect on the pharmacokinetics of albendazole sulfoxide.

Patients with abnormal liver function tests (transaminases) prior to treatment with albendazole should be closely monitored. The treatment should be stopped in case of significant increase in liver enzymes or in case of clinically significant decrease in blood formula numeration (see section 4.4).

Renal failure:

As the elimination of albendazole and its main metabolite, albendazole sulfoxide are negligible, it is unlikely that the clearance of these compounds are modified in patients with renal failure. No dose adaptation of posology is required, however, patients with renal failure should be closely monitored.

Method of administration

Oral route.

Neither purge, nor fast prior treatment is necessary

Some people, notably young children and elderly people might have difficulties swallowing the full tablets. In this case VERZOL oral suspension should be used.

4.3 Contraindications

- Hypersensitivity to albendazole or to any of the components
- Pregnancy and women of childbearing age who do not use an efficient contraceptive method (see section 4.6)
- Breastfeeding

4.4 Special warnings and precautions for use

Neurologic symptoms

A treatment with albendazole might reveal a pre-existing neurocysticercosis, in particular in regions of strong infestation with taeniasis. Patients might feel neurological symptoms such as convulsions, increase in intracranial pressure and focal signs resulting from the inflammatory reactions following the death of the parasite in the brain. Symptoms might appear shortly after the treatment; an adapted treatment with corticoids and anticonvulsants should be immediately started.

Precaution for use when using albendazole for systemic infections (long-term treatment with higher doses):

- Liver disorders

Albendazole might result in a slight to moderate increase in liver transaminases, normalising generally when stopping the treatment. Serious cases of hepatitis have also been reported when treating systemic helminth infections (long-term treatment with higher doses) (see section 4.8). Tests of the liver function should be carried out prior to starting the treatment and at least every second week during the treatment. Albendazole shall be stopped in case of increase in hepatic enzymes (more than twice normal). If reintroducing the treatment is indispensable, this should be done after normalisation of liver enzymes. Moreover, a close

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 31 of 62

monitoring should be carried out, keeping in mind that potential relapses might appear because an allergic mechanism cannot be discarded.

- Medullar depression

Cases of medullar depression have been reported during treatment of systemic helminth infections (long-term treatment with higher doses) (see section 4.8). Numerations of blood formula should be performed when starting the treatment and then after two weeks of treatment with albendazole.

Patients with a liver disease, including liver echinococcosis, seem more likely to develop a medullar depression, leading to pancytopenia, medullar aplasia, agranulocytosis and leucopenia. Then, an increase monitoring of the blood formula is recommended in patients showing a liver disease.

Albendazole shall be stopped in case of significant decrease in the number of blood cells (see section 4.2 and 4.8).

In the treatment of trichinosis, few data are available with albendazole in children under 6 years of age.

In the treatment of trichinosis, because of the activity, in particular on the intestinal forms and of the larvae in the early phase of the tissue migration, it is recommended to administer albendazole as early as possible at the start of the infestation in order to decrease the symptoms and the complications. This treatment remains inactive on the encysted larvae in chronic forms and when it is initiated belatedly.

- Contraception

Before initiating the treatment with albendazole, the doctor should inform the patient of the embryotoxic, teratogenic and aneugenic risks of albendazole, of the necessity of an efficient contraception and of the potential consequences on pregnancy if it occurs during the course of the treatment with albendazole (see section 4.6).

Verzol tablet contains lactose and should not be used by patients with rare hereditary problems of galactose intolerance, lactase deficiency or glucose-galactose malabsorption

Verzol suspension contains sucrose and sorbitol. Patients with rare hereditary problems of fructose intolerance, glucose/galactose malabsorption and sucrase-isomaltase insufficiency should not take this medicine.

The tablets and suspension contain an azo colouring agent (E110) that may cause allergic reactions.

4.5 Interaction with other medicinal products and other forms of interaction

Enzymes inducers anticonvulsivants, ritonavir and rifampicine may have the potential to reduce plasma concentrations of albendazole and of its active metabolite, albendazole sulfoxide with a risk of decrease in its efficacy. Clinical monitoring of the therapeutic efficacy and the potential adaptation of the posology of albendazole during the course of the treatment with an enzymatic inducer and after stopping.

4.6 Fertility, pregnancy and lactation

Female patients

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 32 of 62

Given the aneugenic, embryotoxic and teratogenic potential of albendazole, all the precautions should be taken in order to avoid pregnancy in these female patients. Treatment with albendazole should not be initiated before a negative result to a pregnancy test performed right before the treatment initiation. Women of childbearing age should use an efficient contraceptive method during the treatment and 6 months after stopping the treatment.

Male patients and their female partners

All precaution should be taken in order to avoid pregnancy in the partners of male patients treated with albendazole. It is not known if the presence of albendazole in sperm can cause teratogenic or genotoxic effects on human embryo/foetus. Men or their female partners of childbearing age must be informed of the obligation to use an efficient contraceptive method during all the course of the treatment with albendazole and during 3 months after stopping the treatment. Men whose partners are pregnant should be informed of the obligation to use a condom in order to reduce the exposition of their partner to albendazole.

Pregnancy

Studies in animal showed teratogenic embryotoxic effects in rat and rabbit at doses close to those used in men (see section 5.3). in clinical trials, the data on the use of albendazole during the first term of pregnancy are limited. Albendazole is contraindicated during pregnancy (see section 4.3 and 4.4), especially because there are therapeutical alternatives that are better assessed in terms of safety in pregnant woman. Female patients should be informed of the necessity to consult their doctor immediately in case of pregnancy. This is based on prenatal monitoring targeted on malformations described in animal (skeletal, cranofacial, limbs).

Fertility

In rat or mouse, studies have showed testicular toxicity of albendazole (see section 5.3). albendazole has an aneugic activity, which is a risk factor for alteration of fertility in man.

Breastfeeding

Albendazole is present in human breast milk after a single dose of 400 mg. Because of its aneugenic activity, a risk for the new born child cannot be excluded. In case of a single dose, breastfeeding should be stopped at the time of intake and for at least 5.5 half-lives (about 48 hours) after stopping the treatment. Before initiating breastfeeding, pump all the available breast milk and dispose of it; in case of repeated intakes, breastfeeding is contraindicated.

4.7 Effects on ability to drive and use machines

when driving or using machines, it should be kept in mind that dizziness have been reported after using albendazole (see section 4.8).

4.8 Undesirable effects

The frequency of side effects very common to rare have been determined based on the data from the clinical trials. The frequencies of the other side effects are mainly based on the post-marketing data and are referred to the reported observations rather than the real frequencies. The side effects listed below are classified by organ system and frequency, according to the following convention:

Very common $\geq 1/10$

Common $\geq 1/100$ to $< 1/10$

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 33 of 62

Uncommon $\geq 1/1,000$ to $< 1/100$

Rare $\geq 1/10,000$ to $< 1/1,000$

Very rare $< 1/10,000$

Unknown frequency (cannot be estimated based on the available data).

Intestinal and skin infections (short-term treatment with low doses)

Systemic class organs	Uncommon	Unknown frequency
<i>Immune system disorders</i>		Hypersensitivity reaction, including skin rash, itching and hives
<i>Nervous system disorders</i>	Headaches Dizziness (see section 4.7)	
<i>Gastro-intestinal disorders</i>	Gastro-intestinal symptoms (epigastric or abdominal pains, nausea, vomiting) and diarrhoea	
<i>Hepatobiliary disorders</i>		Increase in liver enzymes (see section 4.4)
<i>Skin and subcutaneous tissue disorders</i>		Polymorphic erythema Stevens-Johnson syndrome

Systemic infections (long-term treatment with higher doses)

Systemic class organs	Very common	Common	Uncommon	Unknown frequency
<i>Haematological and lymph system disorders</i>				<i>Medullar aplasia</i> <i>Leucopenia</i> <i>Pancytopenia</i> <i>Agranulocytosis</i> (see section 4.4)
<i>Immune system disorders</i>			<i>Hypersensitivity reactions including skin rash, itching, hives</i>	
<i>Nervous system disorders</i>	<i>Headaches</i>	<i>Dizziness (see section 4.7)</i>		
<i>Gastro-intestinal disorders</i>		<i>Gastro-intestinal disorders (abdominal pains, nausea and vomiting)</i>		
<i>Hepatobiliary disorders</i>	<i>Slight to moderate increase in liver enzymes</i>		<i>Hepatitis (see section 4.4)</i>	

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 34 of 62

	<i>(see section 4.4)</i>			
<i>Skin and subcutaneous tissue disorders</i>		<i>Reversible alopecia (decrease in thickness of the hair, moderate hair loss)</i>		<i>Polymorphic erythema, Stevens-Johnson syndrome</i>
<i>General disorders and administration site conditions</i>		<i>Fever</i>		

Risk of allergic reactions due to the presence of yellow sunset colouring agent.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

4.9 Overdose

In case of overdose, symptomatic treatment and medical monitoring are recommended.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiparasitics - anthelmintics, ATC code: P02CA03.

Albendazole is a benzimidazole carbamate. Albendazole is broad-spectrum anthelmintics, which is effective against a wide range of intestinal helminths.

Albendazole acts on helminths' cytoskeleton by the inhibition of tubulin polymerisation and thus, their introduction in the microtubules, blocking glucose absorption of parasites and resulting in their death.

Albendazole is also active on *Giardia intestinalis* (or *duodenalis*). It has an irreversible action that is targeted on the ventral disc of the trophozoites by acting on the polymerisation of tubulin and giardine, leading to a disorganisation of the cytoskeleton and micro strips. The ability of adhesion to the enterocytes is decreased, resulting in an inhibition of the growth and multiplication of the parasite.

5.2 Pharmacokinetic properties

Absorption and biotransformation

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 35 of 62

Following the administration, the low proportion of albendazole is absorbed (< 5 %) is metabolised into albendazole sulfoxide and sulfone. The plasma concentration in sulfoxide, the main active circulating metabolite reaches its maximum about two and a half hours after its administration.

The systemic pharmacological effect of albendazole is increased if the dose is administered concomitantly with a fat-rich meal, improving absorption by about 5.

Elimination

The plasma half-life of albendazole sulfoxide is 8 and a half hours.

Albendazole sulfoxide and its metabolites seem to be mainly eliminated by biliary route and for a lower proportion by urinary route.

Specific population

Renal failure: albendazole pharmacokinetics has not been studied in patients with renal failure.

Haptic failure: albendazole pharmacokinetics has not been studied in patients with hepatic failure.

5.3 Preclinical safety data

Degeneration of the seminiferous tubules has been reported in cancerogenesis studies at dose of 100 mg/kg/day in mouse and 20 mg/kg/day in rat. A decrease in the testicle weight has been observed in dog treated with 60 mg/kg/day during 6 months. These doses correspond respectively to 2.4; 0.24 and 2.5 times the maximum therapeutic dose (based on the human equivalence). Albendazole has not altered fertility in males or female rat up to the maximum dose of 30 mg/kg/day, or 0.36 times the maximum therapeutic dose (based on the human equivalence).

Albendazole appeared to be teratogenic and embryotoxic in rat and rabbit.

No cancerogenic potential has been shown during the cancerogenesis studies in rats (20 mg/kg/day) and in mice (400 mg/kg/day). Albendazole did show any genotoxic effects in *in vitro* trials carried out on bacteria and mammal cells cultures, as well as in an *in vivo* micronucleus trial in rodents. A positive result has been reported in another micronucleus study in mouse, and is regarded as resulting from an aneugenic effect of albendazole.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablets:

Maize starch, lactose, sodium methyl parahydroxybenzoate (E219), sodium propyl parahydroxybenzoate (E217), magnesium stearate, purified talc, colloidal anhydrous silica, sunset yellow supra colour (E110).

Suspension:

Saccharose, sorbitol, sodium carboxymethyl cellulose, sunset yellow supra colour (E110), orange flavour, banana flavour, glycerine, sodium methyl parahydroxybenzoate (E219), sodium propyl parahydroxybenzoate (E217), sodium benzoate (E211), sodium saccharin, citric acid, Tween 80, purified water.

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 36 of 62

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Keep out of sight and reach of children.

Protect from heat, light and moisture, store at a temperature not exceeding 30 °C.

6.5 Nature and contents of container

Tablet

Box of 1 tablet in blister pack (PVC-aluminium).

Suspension

Brown-glass bottle of 10 ml suspension.

Box of 20 sticks made of polyethylene terephthalate-aluminium-polyethylene terephthalate-polyethylene of 10 ml.

All packaging might not be marketed.

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Exphar s.a. Zoning Industriel de Nivelles Sud, zone II – Av. Thomas Edison 105 – 1402 Thines (Belgium)

8. CATEGORY OF DISTRIBUTION

Over-the counter medicine

Prescription only medicines

List II

9. MANUFACTURER

Gracure Pharmaceuticals Ltd.,
E-1105, Industrial Area, Phase-III,
Bhiwadi, Dist. Alwar (Raj.) INDIA
Phone +91.11.259.20.748
Fax +91.11.259.20.747

EXPHAR SA	March 2019	VERZOL – Oral suspension
	Administrative Information and Product Information	Module 1 - Page 37 of 62

10. DATE OF REVISION OF THE TEXT

08/2018

1.5.2 Container labelling

Information to appear on the immediate packaging (blister foil)

Name of drug product: present

Pharmaceutical form and Statement of the active substances: present

Name and address of the manufacturer: present

Name and address of the marketing authorization holder (applicant): present

Batch number: present

Manufacturing date: present

Expiry date: present

Information to appear on the outer packaging

Name of drug product: present

Pharmaceutical form and Statement of the active substances: present

Statement “*Read the package leaflet before use.*” present

Special warnings that the drug product must be stored out of the reach and sight of children: present

Storage conditions: present

Name and address of the manufacturer: present

Name and address of the marketing authorization holder (applicant): present

Batch number: present

Manufacturing date: present

Expiry date: present