

SUMMARY OF PRODUCT CHARACTERISTICS GLUCOMET 850 TABLETS
(METFORMIN FILM COATED TABLETS 850MG)

1. Name of the Medicinal Product

Glucomet 850 mg film coated Tablets

2. Qualitative and Quantitative Composition

Each tablet contains 850 mg of Metformin Hydrochloride BP.

White Oblong – shaped, biconvex, film coated tablet embossed ‘COSMOS’ on one side and a ‘BREAKLINE’ on the other side.

3. Pharmaceutical Form

Film coated
Tablet,

4. Clinical Particulars

4.1 Therapeutic Indications

Treatment of type 2 diabetes mellitus, particularly in overweight patients, when dietary management and exercise alone does not result in adequate glycaemic control. In adults, Glucophage may be used as monotherapy or in combination with other oral antidiabetic agents or with insulin. In children from 10 years of age and adolescents, Glucomet may be used as monotherapy or in combination with insulin.

4.2 Posology and Method of administration

Posology

Adults with normal renal function ($GFR \geq 90$ mL/min)

Monotherapy and combination with other oral antidiabetic agents

The usual starting dose is 500 mg or 850 mg metformin hydrochloride 2 or 3 times daily given during or after meals. After 10 to 15 days the dose should be adjusted on the basis of blood glucose measurements. A slow increase of dose may improve gastrointestinal tolerability. The maximum recommended dose of metformin hydrochloride is 3 g daily, taken as 3 divided doses. If transfer from another oral antidiabetic agent is intended: discontinue the other agent and initiate metformin at the dose indicated above.

Combination with insulin

Metformin and insulin may be used in combination therapy to achieve better blood glucose control. Metformin hydrochloride is given at the usual starting dose of 500 mg or 850 mg 2 or 3 times daily, while insulin dosage is adjusted on the basis of blood glucose measurements.

Elderly

Due to the potential for decreased renal function in elderly subjects, the metformin dosage should be adjusted based on renal function. Regular assessment of renal function is necessary.

4.3 Contraindications

Hypersensitivity to metformin or to any of the excipients.

- Diabetic ketoacidosis, diabetic pre-coma.
- Renal failure or renal dysfunction (creatinine clearance < 60 ml/min).
- Acute conditions with the potential to alter renal function such as:
 - Dehydration,
 - Severe infection,
 - Shock,
- Acute or chronic disease which may cause tissue hypoxia such as:
 - Cardiac or respiratory failure,
 - Recent myocardial infarction,
 - Shock
- Hepatic insufficiency, acute alcohol intoxication, alcoholism

4.4 Special warnings and precautions for use

Lactic acidosis:

Lactic acidosis is a rare, but serious (high mortality in the absence of prompt treatment), metabolic complication that can occur due to metformin accumulation.

Diagnosis:

The risk of lactic acidosis must be considered in the event of non-specific signs such as muscle cramps with digestive disorders as abdominal pain and severe

asthenia. This can be followed by acidotic dyspnea, abdominal pain, hypothermia and coma.

Renal function:

As metformin is excreted by the kidney, creatinine clearance (this can be estimated from serum creatinine levels using the Cockcroft-Gault formula) should be determined before initiating treatment and regularly thereafter:

- at least annually in patients with normal renal function,
- at least two to four times a year in patients with creatinine clearance levels at the limit of normal and in elderly subjects.

Administration of iodinated contrast media:

The intravascular administration of iodinated contrast media in radiological studies can lead to renal failure. This may lead to metformin accumulation and risk of lactic acidosis. Metformin must be discontinued prior to, or at the time of the test and not reinstituted until 48 hours afterwards, and only after renal function has been re-evaluated and found to be normal.

Surgery:

Metformin should be discontinued 48 hours before elective surgery with general spinal or peridural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition provided normal renal function has been established.

Other precautions:

All patients should continue their diet with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.

The usual laboratory tests for diabetes monitoring should be performed regularly.

Metformin alone never causes hypoglycaemia, although caution is advised when it is used in combination with insulin or other oral antidiabetics (e.g. sulphonylureas or meglitinides).

The tablet shells may be present in the faeces. Patients should be advised that this is normal.

4.5 Interaction with other medicinal products and other forms of interaction

- Concomitant use not recommended

Alcohol

Acute alcohol intoxication is associated with an increased risk of lactic acidosis in acute alcohol intoxication, particularly in case of:

- fasting or malnutrition,
- hepatic insufficiency.

Avoid consumption of alcohol and alcohol-containing medications.

Iodinated contrast media

Intravascular administration of iodinated contrast media may lead to renal failure, resulting in metformin accumulation and a risk of lactic acidosis.

Metformin hydrochloride must be discontinued prior to, or at the time of the test and not reinstituted until 48 hours afterwards, and only after renal function has been re-evaluated and found to be normal.

- Combinations requiring precautions for use

Medicinal products with intrinsic hyperglycaemic activity (e.g. glucocorticoids (systemic and local routes) and sympathomimetics). More frequent blood glucose monitoring may be required, especially at the beginning of treatment. If necessary, adjust the metformin dosage during therapy with the other drug and upon its discontinuation.

- Diuretics, especially loop diuretics
- They may increase the risk of lactic acidosis due to their potential to decrease renal function

4.6 Pregnancy and lactation

Pregnancy

Uncontrolled diabetes during pregnancy (gestational or permanent) is associated with increased risk of congenital abnormalities and perinatal mortality.

A limited amount of data from the use of metformin in pregnant women does not indicate an increased risk of congenital abnormalities. Animal studies do not indicate harmful effects with respect to pregnancy, embryonic or fetal development, parturition or postnatal development.

Lactation

Metformin is excreted into human breast milk. No adverse effects were observed in breastfed newborns/infants. However, as only limited data are available, breastfeeding is not recommended during metformin treatment. A decision on whether to discontinue breast-feeding should be made, taking into account the benefit of breast-feeding and the potential risk to adverse effect on the child.

Fertility

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

4.7 Effects on ability to drive and use machines

Metformin monotherapy does not cause hypoglycaemia and therefore has no effect on the ability to drive or to use machines.

However, patients should be alerted to the risk of hypoglycaemia when metformin is used in combination with other antidiabetic agents (e.g. sulphonylureas, insulin, or meglinitides).

4.8 Undesirable Effects

During treatment initiation, the most common adverse reactions are nausea, vomiting, diarrhoea, abdominal pain and loss of appetite, which resolve spontaneously in most cases.

The following adverse reactions may occur with Glucomet 850.

Frequencies are defined as follows: very common: $>1/10$; common $>1/100$, $<1/10$; uncommon $>1/1,000$, $<1/100$; rare $>1/10,000$, $<1/1,000$; very rare $<1/10,000$.

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Metabolism and nutrition disorders

Very rare: Lactic acidosis.

Decrease of vitamin B12 absorption with decrease of serum levels during long-term use of metformin. Consideration of such an aetiology is recommended if a patient presents with megaloblastic anaemia.

Nervous system disorders

Common: Taste disturbance

Gastrointestinal disorders

Very common:

Gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite. These undesirable effects occur most frequently during initiation of therapy and resolve spontaneously in most cases. A slow increase of the dose may also improve gastrointestinal tolerability.

Hepatobiliary disorders

Isolated reports: Liver function tests abnormalities or hepatitis resolving upon metformin discontinuation.

Skin and subcutaneous tissue disorders

Very rare: Skin reactions such as erythema, pruritus, urticaria

5 Overdose

Hypoglycaemia has not been seen with metformin doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose or concomitant risks of metformin may lead to lactic acidosis. Lactic acidosis is a medical emergency and must be treated in hospital. The most effective method to remove lactate and metformin is haemodialysis.

6 Pharmacological Properties

6.1 Pharmacodynamic Properties

Metformin is an antihyperglycemic agent which improves glucose tolerance in patients with type 2 diabetes, lowering both basal and postprandial plasma glucose. Its pharmacologic mechanisms of action are different from other classes of oral antihyperglycemic agents. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. Unlike sulfonylureas, metformin does not produce hypoglycemia in either patients with type 2 diabetes or normal subjects (except in special circumstances) and does not cause hyperinsulinemia. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may actually decrease.

6.2 Pharmacokinetic Properties

Absorption

After an oral dose of the prolonged release tablet, metformin absorption is significantly delayed compared to the immediate release tablet with a T_{max} at 7 hours (T_{max} for the immediate release tablet is 2.5 hours).

At steady state, similar to the immediate release formulation, C_{max} and AUC are not proportionally increased to the administered dose. The AUC after a single oral administration of 2000mg of metformin prolonged release tablets is similar to that observed after administration of 1000mg of metformin immediate release tablets b.i.d.

Intrasubject variability of C_{max} and AUC of metformin prolonged release is comparable to that observed with metformin immediate release tablets.

When the prolonged release tablet is administered in fasting conditions the AUC is decreased by 30% (both C_{max} and T_{max} are unaffected).

Mean metformin absorption from the prolonged release formulation is almost not altered by meal composition.

Distribution

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean Vd ranged between 63-276 L.

Metabolism

Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

Elimination

Renal clearance of metformin is > 400 ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours. When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

6.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies on safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity reproduction.

7 Pharmaceutical Particulars

7.1 List of Excipients

Povidone K-30BP

Maize starch BP

Sodium starch glycollate BP

Isopropyl alcohol BP

Colloidal Anhydrous Silica BP

Magnesium Stearate BP

COATING

Hypromellose BP

Propylene glycol BP

Purified talc BP

Titanium dioxide BP

Isopropyl alcohol BP

Purified water BP

POLISHING

Carnauba wax BP

Hard paraffine BP

Dichlorométhane BP

7.2 Incompatibilities

None

7.3 Shelf life

3 Years

7.4 Special precautions for storage

Store in a dry place below 30°C.

7.5 Nature and contents of container

PVC/ALU blister packing

7.6 Instructions for use, handling and disposal

No special requirements

8 Registrant

Cosmos Limited

9 Manufacturer

Cosmos Limited

Rangwe Road; Off Lunga Lungu, Industrial Area

P.O Box 41433, GPO 00100-Nairobi

Kenya

Telephone: + (254-20) 2519603/4/5, 020 8042200/2/3/4/5

Telefax: +254-020-8096280/1

E-mail: admin@cosmos-pharm.com