

TRIXONE
CEFTRIAXONE SODIUM INJECTION
B.P. 250MG/500MG/1.0G

COMPOSITION:

Each vial contains:
Ceftriaxone Sodium BP equivalent to Ceftriaxone 250mg/500mg/1g

INDICATIONS:

Meningitis, Septicaemia, Typhoid, UTI, Prophylaxis in surgical infections

DOSAGE:

MENINGITIS: Adults: 4gm i.v. Initially followed by 2 gm i.v. once daily for 10 days or 72 hours after afebrile.

Children: 75-100 mg/kg wt. for 7-9 days.

TYPHOID: Adults: 4gm daily for 2 days followed by 2 gm daily for 2 days.

Children: 75 mg/kg body wt daily for 7 days.

Other conditions: Adults: 1-2 gms daily

Children: 50-75 mg/kg body wt. daily.

DIRECTION FOR USE:

For 1 gm: Dissolve the content of vial in 9.6 ml of sterile water for Injection BP for IV administration and 3.6 ml for IM administration.

For 500 mg: Dissolve the content of vial in 4.8 ml of sterile water for Injection BP for IV administration and 0.9 ml for IM administration.

For 250 mg: Dissolve the content of vial in 2.4 ml of sterile water for Injection BP for IV administration and 0.9 ml for IM administration.

The reconstituted solution should be used immediately after preparation.

CONTRA-INDICATIONS:

Hypersensitivity

SPECIAL PRECAUTIONS:

Hepatic or renal dysfunction in Neonates.

Paediatrics: Avoid in premature neonates

Pregnancy and Lactation: Use with caution.

SIDE-EFFECTS:

Diarrhoea, hypo-prothrombinaemia, pseudomembranous colitis.

DRUG INTERACTIONS:

Disulfiram like reaction with alcohol.

STORAGE:

KEEP OUT OF REACH OF CHILDREN

Store below 25°C. in a dry place. Protect from light.

SHELF-LIFE:

Three years.

PRESENTATION:

250 mg, 500 mg and 1 gm.

Manufactured in China by:

Reyoung Pharmaceutical Co., Ltd.

No.1 Ruiyang Road, Yiyuan County,

Shandong Province, P.R. China.



ABACUS

Manufactured for:

Abacus Pharma (A) Limited

EAST AFRICA

120*170mm (机装小盒说明书尺寸)