

treatments have been observed in the clinical trial programme.

PHARMACEUTICAL PARTICULARS:

INCOMPATIBILITY

Not applicable.

STORAGE CONDITION

Store below 30°C, Protect from moisture.

NATURE AND CONTENTS OF CONTAINER

10 tablets in Alu-Alu blister pack, 3 such blisters in a printed carton along with Pack Insert.

MANUFACTURING AUTHORISATION HOLDER AND MANUFACTURER

Manufacturing Authorisation Holder	Manufacturer
Ajanta Pharma Limited Ajanta House, Charkop Kandivli (West) Mumbai - 400 067 India. Tel : +91-22-6606 1000 Fax : (0091) 22-6606 1200/ 300 Email : info@ajantapharma.com	Ajanta Pharma Limited Z/103/A, Dahaj Sez II, Bhanuch -392 130 India. Tel : 91-22-6606 1000 Fax : (0091) 22-66061200/ 300 Email : info@ajantapharma.com

Version No.: 00

Last Revision Date: May, 2021

VILDARIL M

Patient Information Leaflet

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

What is in this leaflet

1. What Vildaril M is and what it is used for

2. What you need to know before you take Vildaril M

3. How to take Vildaril M

4. Possible side effects

5. How to store Vildaril M

6. Contents of the pack and other information

1. What Vildaril M is and what it is used for

The active substances of Vildaril M, vildagliptin and metformin, belong to a group of medicines called "oral anti-diabetics".

Vildaril M is used to treat adult patients with type 2 diabetes. This type of diabetes is also known as noninsulin-dependent diabetes mellitus.

Type 2 diabetes develops if the body does not make enough insulin or if the insulin that the body makes does not work as well as it should. It can also develop if the body produces too much glucagon.

Both insulin and glucagon are made in the pancreas. Insulin helps to lower the level of sugar in the blood, especially after meals. Glucagon triggers the liver to make sugar, causing the blood sugar level to rise.

How Vildaril M works

Both active substances, vildagliptin and metformin, help to control the level of sugar in the blood. The substance vildagliptin works by making the pancreas produce more insulin and less glucagon. The substance metformin works by helping the body to make better use of insulin. This medicine has been shown to reduce blood sugar, which may help to prevent complications from your diabetes.

2. What you need to know before you take Vildaril M

Do not take Vildaril M

- if you are allergic to vildagliptin, metformin or any of the other ingredients of this medicine. If you think you may be allergic to any of these, talk to your doctor before taking Vildaril M.
- if you have uncontrolled diabetes, with, for example, severe hyperglycaemia (high blood glucose), nausea, vomiting, diarrhoea, rapid weight loss, lactic acidosis (see "Risk of lactic acidosis" below) or ketoacidosis. Ketoacidosis is a condition in which substances called ketone bodies accumulate in the blood and which can lead to diabetic pre-coma. Symptoms include stomach pain, fast and deep breathing, sleepiness or your breath developing an unusual fruity smell.
- if you have recently had a heart attack or if you have heart failure or serious problems with your blood circulation or difficulties in breathing which could be a sign of heart problems.
- if you have severely reduced kidney function.
- if you have a severe infection or are seriously dehydrated (have lost a lot of water from your body).
- if you are going to have a contrast x-ray (a specific type of x-ray involving an injectable dye).

Please also see information about this in section "Warnings and precautions".

- if you have liver problems.
- if you drink alcohol excessively (whether every day or only from time to time).

- if you are breast-feeding (see also "Pregnancy and breast-feeding").

Warnings and precautions

Risk of lactic acidosis
Vildaril M may cause a very rare, but very serious side effect called lactic acidosis, particularly if your kidneys are not working properly. The risk of developing lactic acidosis is also increased with uncontrolled diabetes, serious infections, prolonged fasting or alcohol intake, dehydration (see further information below), liver problems and any medical conditions in which a part of the body has a reduced supply of oxygen (such as acute severe heart disease).

If any of the above apply to you, talk to your doctor for further instructions.

Stop taking Vildaril M for a short time if you have a condition that may be associated with dehydration (significant loss of body fluids) such as severe vomiting, diarrhoea, fever, exposure to heat or if you drink less fluid than normal. Talk to your doctor for further instructions.

Stop taking Vildaril M and contact a doctor or the nearest hospital immediately if you experience some of the symptoms of lactic acidosis, as this condition may lead to coma.

Symptoms of lactic acidosis include:

- vomiting
- stomach ache (abdominal pain)
- muscle cramps
- a general feeling of not being well with severe tiredness
- difficulty in breathing
- reduced body temperature and heartbeat

Lactic acidosis is a medical emergency and must be treated in a hospital.

Vildaril M is not a substitute for insulin. Therefore, you should not receive Vildaril M for the treatment of type 1 diabetes.

Talk to your doctor, pharmacist or nurse before taking Vildaril M if you have or have had a disease of the pancreas.

Talk to your doctor, pharmacist or nurse before taking Vildaril M if you are taking an anti-diabetic medicine known as a sulphonylurea. Your doctor may want to reduce your dose of the sulphonylurea when you take it together with Vildaril M in order to avoid low blood glucose (hypoglycaemia).

If you have previously taken vildagliptin but had to stop taking it because of liver disease, you should not take this medicine.

Diabetic skin lesions are a common complication of diabetes. You are advised to follow the recommendations for skin and foot care that you are given by your doctor or nurse. You are also advised to pay particular attention to new onset of blisters or ulcers while taking Vildaril M. Should these occur, you should promptly consult your doctor.

If you need to have major surgery you must stop taking Vildaril M during and for some time after the procedure. Your doctor will decide when you must stop and when to restart your treatment with Vildaril M.

A test to determine your liver function will be performed before the start of Vildaril M treatment, at three-month intervals for the first year and periodically thereafter. This is so that signs of increased liver enzymes can be detected as early as possible.

During treatment with Vildaril M, your doctor will check your kidney function at least once a year or more frequently if you are elderly and/or have worsening renal function.

Your doctor will test your blood and urine for sugar regularly.

Children and adolescents
The use of Vildaril M in children and adolescents up to 18 years of age is not recommended.

Other medicines and Vildaril M
If you need to have an injection of a contrast medium that contains iodine into your bloodstream, for example in the context of an X-ray or scan, you must stop taking Vildaril M before or at the time of the injection. Your doctor will decide when you must stop and when to restart your treatment with Vildaril M.

Tell your doctor if you are taking, have recently taken or might take any other medicines. You may need more frequent blood glucose and kidney function tests, or your doctor may need to adjust the dosage of Vildaril M. It is especially important to mention the following:

- glucocorticoids generally used to treat inflammation

- beta-2 agonists generally used to treat respiratory disorders
- other medicines used to treat diabetes
- medicines which increase urine production (diuretics)

• medicines used to treat pain and inflammation (NSAID and COX-2-inhibitors, such as ibuprofen and celecoxib)

• certain medicines for the treatment of high blood pressure (ACE inhibitors and angiotensin II receptor antagonists)

• certain medicines affecting the thyroid, or

• certain medicines affecting the nervous system.

Vildaril M with alcohol
Avoid excessive alcohol intake while taking Vildaril M since this may increase the risk of lactic acidosis (please see section "Warnings and precautions").

Pregnancy and breast-feeding
• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

• If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will discuss with you the potential risk of taking Vildaril M during pregnancy.

difficult breathing, sudden onset of rash or hives, which may indicate a reaction called "angioedema".

Liver disease (hepatitis) (rare). Symptoms include yellow skin and eyes, nausea, loss of appetite/ dark-coloured urine, which may indicate liver disease (hepatitis).

Inflammation of the pancreas (pancreatitis) (frequency not known). Symptoms include severe and persistent pain in the abdomen (stomach area), which might reach through to your back, as well as nausea and vomiting.

Other side effects
Some patients have experienced the following side effects while taking Vildaril M:

• Very common (may affect more than 1 in 10 people): nausea, vomiting, diarrhoea, pain in and around the stomach (abdominal pain), loss of appetite.

• Common (may affect up to 1 in 10 people): dizziness, headache, trembling that cannot be controlled, metallic taste, low blood glucose.

• Uncommon (may affect up to 1 in 100 people): joint pain, tiredness, constipation, swollen hands, ankle or feet (oedema).

• Very rare (may affect up to 1 in 10,000 people): sore throat, runny nose, fever; signs of a high level of lactic acid in the blood (known as lactic acidosis) such as drowsiness or dizziness, severe nausea or vomiting, abdominal pain, irregular heart beat or deep, rapid breathing, redness of the skin, itching; decreased vitamin B12 levels (paleness, tiredness, mental symptoms such as confusion or memory disturbances).

Some patients have experienced the following side effects while taking Vildaril M and a sulphonylurea:

• Common: dizziness, tremor, weakness, low blood glucose, excessive sweating.

Some patients have had the following side effects while taking Vildaril M and insulin:

• Common: headache, chills, nausea (feeling sick), low blood glucose, heartburn.

• Uncommon: diarrhoea, flatulence.

Since this product has been marketed, the following side effects have also been reported:

• Frequency not known (cannot be estimated from the available data): itchy rash, inflammation of the pancreas, localised peeling of skin or blisters, muscle pain.

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Vildaril M

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the blister and carton after "EXP". The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Store in the original package (blister) in order to protect from moisture

6. Contents of the pack and other information

What Vildaril M 50/500, Vildaril M 50/850 and Vildaril M 50/1000 Contains:

List of Excipients
VILDARIL M 50/500(Vildagliptin 50 mg and Metformin Hydrochloride 500 mg Tablets)

Lactose (Monohydrate), Hydroxypropyl cellulose, Isopropyl Alcohol, Sodium Starch Glycolate Type B, Microcrystalline Cellulose, Hydrophobic Colloidal Silica, Magnesium Stearate, InstaMoistshield A21R01476 Yellow (Hypromellose, Ethyl Cellulose, Tracelac, Talc, Titanium Dioxide, Yellow iron oxide, Red iron oxide) and Dichloromethane

VILDARIL M 50/850(Vildagliptin 50 mg and Metformin Hydrochloride 850 mg Tablets)

Lactose (Monohydrate), Hydroxypropyl cellulose, Isopropyl Alcohol, Sodium Starch Glycolate Type B, Microcrystalline Cellulose, Hydrophobic Colloidal Silica, Magnesium Stearate, InstaMoistshield A21R01476 Yellow (Hypromellose, Ethyl Cellulose, Tracelac, Talc, Titanium Dioxide, Yellow iron oxide, Red iron oxide) and Dichloromethane

VILDARIL M 50/1000(Vildagliptin 50 mg and Metformin Hydrochloride 1000 mg Tablets)

Lactose (Monohydrate), Hydroxypropyl cellulose, Isopropyl Alcohol, Sodium Starch Glycolate Type B, Microcrystalline Cellulose, Hydrophobic Colloidal Silica, Magnesium Stearate, InstaMoistshield A21R01476 Yellow (Hypromellose, Ethyl Cellulose, Tracelac, Talc, Titanium Dioxide, Yellow iron oxide, Red iron oxide) and Dichloromethane

InstaMoistshield A21R01474 Yellow (Hypromellose, Ethyl Cellulose, Tracelac, Talc, Titanium Dioxide, Yellow iron oxide, Red iron oxide) and Dichloromethane

VILDARIL M 50/1000(Vildagliptin 50 mg and Metformin Hydrochloride 1000 mg Tablets)

Lactose (Monohydrate), Hydroxypropyl cellulose, Isopropyl Alcohol, Sodium Starch Glycolate Type B, Microcrystalline Cellulose, Hydrophobic Colloidal Silica, Magnesium Stearate, InstaMoistshield A21R01475 Yellow (Hypromellose, Ethyl Cellulose, Tracelac, Talc, Titanium Dioxide, Yellow iron oxide, Red iron oxide) and Dichloromethane

What Vildaril M 50/500, Vildaril M 50/850 and Vildaril M 50/1000 look and contents of the pack.

VILDARIL M 50/500 (Vildagliptin 50 mg and Metformin Hydrochloride 500 mg Tablets)

Beige to creamish yellow coloured, oval shaped, film coated tablets, plain on both sides.

VILDARIL M 50/850 (Vildagliptin 50 mg and Metformin Hydrochloride 850 mg Tablets)

Yellow to dark yellow coloured, oval shaped, film coated tablets, plain on both sides.

VILDARIL M 50/1000 (Vildagliptin 50 mg and Metformin Hydrochloride 1000 mg Tablets)

Dark yellow to brownish yellow coloured, oval shaped, film coated tablets, plain on both sides

10 tablets in Alu-Alu blister pack, 3 such blisters in a printed carton along with Pack Insert.

MANUFACTURING AUTHORISATION HOLDER AND MANUFACTURER

Manufacturing Authorisation Holder	Manufacturer
Ajanta Pharma Limited Ajanta House, Charkop Kandivli (West) Mumbai - 400 067 India. Tel : +91-22-6606 1000 Fax : (0091) 22-6606 1200/ 300 Email : info@ajantapharma.com	Ajanta Pharma Limited Z/103/A, Dahaj Sez II, Bhanuch -392 130 India. Tel : 91-22-6606 1000 Fax : (0091) 22-66061200/ 300 Email : info@ajantapharma.com

For any information about this medicinal product, please contact the manufacturing authorisation holder.

Date of Publication or Revision: May, 2021