



Regd. Office & Factory : Plot No. 33, Sector II, The Vasai Taluka Industrial Co-op. Estate Ltd. Gaurai-pada, Vasai (E), Dist. Thane - 401 208, INDIA.
Tel. : 95250 - 2455801 / 2452714 / 2453525 • Fax : 95250 - 2452074 (0091 - 250 - 2452074) • Email : agog@vsnl.net & agogpharma@rediffmail.com

Leaflet

Package leaflet: Information for the user
AGOVIT-6 TABLETS
(Pyridoxine Tablet BP 25 mg)

Read all of this leaflet carefully before taking this medicine

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your health care provider or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptom are the same as yours.
- If you get any side effect, talk to your doctor or pharmacist. This includes any possible side effect not listed in this leaflet.

What is in this leaflet?

1. What Agovit-6 Tablets are and what they are used for?
2. What do you need to know before you take Agovit-6 Tablets?
3. How to take Agovit-6 Tablets?
4. Possible Side Effects
5. How to store Agovit-6 Tablets?
6. Further information

1. What Agovit-6 Tablets are and what they are used for?

Agovit-6 tablets contain the active substance Pyridoxine Hydrochloride. Pyridoxine is also called as Vitamin B₆. Pyridoxine tablets are used to treat and prevent vitamin B₆ deficiency in the body. Vitamin B₆ helps the body to use and store energy from protein and carbohydrates in food. It also helps the body form hemoglobin, the substance in red blood cells that carries oxygen around the body.

Pyridoxine tablets are used:

- To treat Vitamin B₆ deficiency
- To treat and prevent Peripheral Neuropathy (damage to the nerves) caused by Isoniazid (a tuberculosis medicine)
- Sideroblastic Anaemia

2. What you need to know before you take Agovit-6 Tablets?

Do not take this medicine if

- You are allergic (hypersensitive) to Pyridoxine Hydrochloride

Take special care with Agovit-6 Tablets

If you have

- Severe allergies or Asthma
- Liver or kidney disease
- Diabetes
- Have a condition called Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency
- Been taking any of these medicine listed in "taking with other medicines"

Taking with other medicines

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without prescription.

Tell your doctor if you are taking any of these following medicines

- Alletamine (to treat ovarian cancer)
- Isoniazid, Cycloserine (to treat tuberculosis)
- Levodopa (to treat Parkinson's disease)
- Azithromycin, Clarithromycin, Erythromycin (to treat bacterial infections)
- Phenytoin, Phenobarbital, Primidone (to treat epilepsy)
- Cisplatin (to treat bladder cancer, testicular cancer and ovarian cancer)
- Penicillamine (to treat arthritis)
- Hydralazine (to treat high blood pressure)

Taking Agovit-6 Tablets with food and drink

Talk to your doctor or pharmacist for best advice on foods and drinks to avoid while taking Agovit-6 tablets. Do not consume alcohol while taking this medicine as it can make you feel more drowsy and sleepy.

Pregnancy and breast-feeding

Talk to your doctor or pharmacist before taking this medicine if you are pregnant or breast-feeding, think you might be pregnant or are planning to have a baby.

Driving and using machine

Agovit-6 tablets do not affect the ability to drive or operate machinery.

3. How to take Agovit-6 Tablets?

Always take this medicine exactly as your doctor or health care provider has told you. You should check with your doctor or health care provider if you are not sure.

Unless otherwise prescribed by the doctor, the recommended doses are:

Adults:

Vitamin B6 Deficiency: 2.5-10 mg/day orally

Peripheral Neuropathy: Patients treated with isoniazid for Mycobacterium tuberculosis: 25-50 mg/day oral

If you take more Agovit-6 Tablets than you should

If you have accidentally taken more tablets in one day than your doctor has told you to, contact your doctor or go to nearest hospital emergency department immediately.

If you forget to take Agovit-6 Tablets

Do not take a double dose to make up for a forgotten dose if you forget to take these tablets.

If you stop taking Agovit-6 Tablets

Do not stop taking your tablets, even if you feel better. It is important to take the tablets as long as the doctor has told you to, otherwise your medicine might not work properly.

If you have any further questions on the use of this product, ask your doctor or health care provider.

Agovit-6 Tablets contains Lactose

Lactose is a type of sugar. If you have been told by a doctor that you cannot tolerate sugar, talk to your doctor or healthcare provider before taking this medicine.

4. Possible side effect:

Like all medicines, Pyridoxine can cause side effects, although not everybody gets them.

Very common side effect: Rashes and Headache

Common side effect: Dry Mouth, Feeling Tired or Dizzy.

Uncommon side effect: Vomiting and Liver Injury.

Rare side effect: Swollen Tongue, Mouth, Lips and Tingling, Pricking or Numbness of Skin.

If any of the side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or health care provider.

5. How to store Agovit-6 Tablets

Do not store above 30°C. Protect from light.

Keep out of the reach and sight of children.

Store in the original carton.

Do not use this medicine after the expiry date stated on the carton (MM/YYYY)

6. Further Information

What Agovit-6 Tablets contain:

Each uncoated tablet contains:

Pyridoxine Hydrochloride BP 25 mg

Excipients q.s.

This product contains "Lactose".

What Agovit-6 tablets looks like and contents of the pack

White, circular, flat uncoated tablet having a breakline on one side and other side is plain of each tablet.

Shelf life: 36 months

Packing/pack size: Blister pack of 10 x 10 tablets

Jar Pack of 1000 tablets

This leaflet is last revised on July 2024

Marketing Authorisation Holder and Manufacturer



AGOG PHARMA LTD.

Plot No. 33, Sector - II, The Vasai Taluka Indl.

Co-op. Estate Ltd., Vasai (E), Dist. Thane, INDIA.

110 mm

1105/02/01

70 mm